

Utility of a Handheld Blood Ketone Meter as a Postmortem Indicator of Diabetic Ketoacidosis

Richard E. Seeber II, BA, BS

Daniel S. Atherton, MD

Brandi McCleskey, MD



Outline

- Research question
- Biochemistry review: ketone bodies; DKA
- Methods
- Present data collected at the Jefferson County OCME
- Discussion of findings
- Two CME/SAMs questions



Research questions

- Can a commercially available blood ketone meter reliably measure post-mortem blood ketone levels such that an accurate diagnosis of antemortem DKA can be made?
- What implications would such a test have for medical examiners' and coroners' offices?



Biochemistry: Glucose and ketones 1

- During life, blood glucose levels may be reliably tested using commercially-available blood glucose meters; in a living patient with DKA, blood glucose levels may be significantly deranged (>250 mg/dL)
 - ♦ Postmortem blood glucose changes



LAB MEDICINE
Knowledge that will change your world


Biochemistry: Glucose and ketones 2

- In life, blood solutes equilibrate with the vitreous humor.
 - ♦ Postmortem vitreous glucose stability
 - ♦ “Gold standard” for postmortem DKA diagnosis when taking clinical picture, PMH into consideration.
 - Testing may be technically challenging or expensive, particularly in smaller or rural offices

LAB MEDICINE
Knowledge that will change your world

Biochemistry: Glucose and ketones 3

- During times of starvation, nutritional deficiency, or “pseudostarvation” (e.g. uncontrolled diabetes mellitus,) the body produces metabolic fuel by catabolism:
 - ♦ Three ketone bodies:



LAB MEDICINE
Knowledge that will change your world

Biochemistry: Glucose and ketones 4

- Postmortem blood ketone values, in contrast with blood glucose values, remain relatively stable [source]
 - ♦ Blood glucose meters; blood ketone meters for *in vivo* DKA screening
 - Blood ketone meters analyze blood concentration of **beta-hydroxybutyrate**, the most prevalent ketone in the body in setting of DKA.
 - Per packaged instructions, ketone levels **>1.5 mmol/L are concerning for DKA.**





Methods


- Brief retrospective pilot study; gas chromatography detected blood acetone in selected decedents ⑦ post-mortem blood assayed for ketones
- Follow-up: Controls and suspected DKA cases were prospectively chosen; decedents' blood samples were analyzed for ketones (screen) and vitreous samples for glucose (confirmatory).





Causes of death in retrospective pilot











Conclusions/Discussion

- Retrospective pilot study results – Post-mortem ketone testing is sensitive *but not specific* for DKA for retrospectively collected samples; vitreous remains most specific gold standard.
- Prospective study: Post-mortem ketone testing is sensitive and, in this small sample, specific for DKA. All decedents dying of DKA had values > 2.6 mmol/L; all others had values < 1 mmol/L (ULN: 1.5 mmol/L)
- Ramifications for OC/Mes
- The art of post-mortem diagnosis

LAB MEDICINE
Knowledge that will change your world

CME Question 1

Question 1: Which of the following is the most prevalent ketone body in the blood in the setting of acute diabetic ketoacidosis?

Answer A: Acetone

- Answer B: β -Ketopentanoate
- Answer C: Acetoacetate
- Answer D: Dihydropteroate
- Answer E: β -Hydroxybutyrate

LAB MEDICINE
Knowledge that will change your world

CME Question 2

- Question 2: Which of the following blood ketone values warrants further work-up for diabetic ketoacidosis?
- Answer A: 0 mmol/L
- Answer B: 0.06 - 0.6 mmol/L
- Answer C: 0.6 - 1.0 mmol/L
- Answer D: 1.0 - 1.5 mmol/L
- Answer E: Greater than 1.5 mmol/L



CME Question references

1. Laffel L. Ketone bodies: a review of physiology, pathophysiology and application of monitoring to diabetes. Diabetes/Metabolism Research and Reviews 1999; 15:412.
2. Charles RA, Bee YM, Eng PHK, et al. Point-of-care blood ketone testing: screening for diabetic ketoacidosis at the emergency department. Singapore Medical Journal 2007; 48:986.



Questions?